

## EFFECTS OF WATER PHYSICOCHEMICAL VARIATIONS ON THE HATCHABILITY AND SURVIVAL OF *Clarias gariepinus* (BURCHELL, 1822) LARVAE

AWOKE, J. S.<sup>1,2</sup>, NWAKPA, J. N.<sup>1</sup>, OTI, E. E.<sup>1</sup>

<sup>1</sup>Department of Fisheries and Aquaculture, Ebonyi State University, P.M.B. 053, Abakaliki - Nigeria

<sup>2</sup>Department of Biology, Ebonyi State College of Education, Ikwo, P.M.B. 004, Abakaliki, Nigeria.

Corresponding author: Awoke, J.S.

.Phone no: 07060517022

E-mail: [judeawoke786@yahoo.com](mailto:judeawoke786@yahoo.com).

### ABSTRACT

*This study investigated the effects of physicochemical parameters of four different water sources on hatchability and survival of Clarias gariepinus larvae. The water sources used were: rain water (T1), borehole water (T2), river water (T3) and well water (T4). The study entailed artificial induced breeding of C. gariepinus using the aforementioned water sources as incubation and culture medium of the hatched larvae for 14 days. Vital water physicochemical parameters assessed were temperature, pH, dissolved oxygen and ammonia, conductivity and total hardness. Hatchability rate was estimated 24 hours after incubation whereas larval survival rate was determined at 14 days' post hatching. Data from the experiment were subjected to one-way analysis of variance (ANOVA) at ( $P < 0.05$ ) significance level. At the end of the experiment, results obtained for water physicochemical parameters in the four water treatments for conductivity and total hardness were significantly different ( $P < 0.05$ ). Rain water (T1) recorded a low mean conductivity value of  $24.3 \pm 0.88 \mu S^{cm}$  and low mean total hardness value of  $3.33 \pm 0.3 mg^l$ . Percentage hatchability in all treatments was relatively high with the highest mean value of  $93.1 \pm 1.60$  % obtained in T1. Larval survival rate at 14 days post hatching was comparatively above average with the highest mean value of  $63.7 \pm 2.41$  % recorded in T2 while T1 recorded the lowest mean value of  $44.2 \pm 0.55$  %. This study has shown that borehole water (T2) is the most suitable water source for hatching and survival of C. gariepinus larvae.*

**Keywords:** Water quality, Hatching, Artificial propagation, Fish seed, Water hardness

### INTRODUCTION

Physicochemical parameters of water are extremely essential to the growth and survival of organisms that inhabit the water bodies. They play important roles in the physiological development, reproduction and general well-being of aquatic organisms such

as fish (Ukwe and Abu, 2016). In aquaculture, it is always an excellent practice to use a reliable source of water during breeding operations. This is because physicochemical characteristics of water are known to influence fish breeding performance in positive and negative ways

(Akombo *et al.*, 2018). Hence, the success of every fish breeding venture is reliant on the quality of water on hand in the hatchery. Fertilization, hatching and early survival of larvae are crucial for successful propagation of the African catfishes. Optimum hatchability of eggs and survival of hatchlings are equally dependent on good water quality (Ataguba *et al.*, 2009).

A number of hatcheries are challenged with some operational difficulties such as low hatchability and low survival rates (Kareem *et al.*, 2017). These drawbacks are linked to water quality in the hatchery system (Akinwale *et al.*, 2006). According to Uka and Kalu, (2019) water quality is the physical, chemical and biological constituents of water. A good water quality for fish culture ought to satisfy the minimum conditions for reproduction, survival and growth of cultivable fish species. Optimal fish production to a very large extent is absolutely dependent on the physical, chemical and biological qualities of water (Bhatnagar and Devi, 2019). Hence, successful hatchery operations require an understanding of water quality. However, it is apparent that, poor water quality in hatcheries may cause low survival of eggs, larvae or fry (Summerfelt, 1998).

In Ebonyi State, South Eastern Nigeria there has been reports of breeding failures and high larval mortalities during breeding operations (Ebiriekwe, 2012). This crisis of larval mortalities during early stages of breeding had been documented, with water quality being the major causative factor (Awoke *et al.*, 2021). According to Opoke and Osayande, (2018) the quality of surface and ground waters in Abakaliki metropolis has been compromised due to its hardness. This has been attributed to the occurrence of mineral deposits and heavy metal ores under the ground (Nwabunike, 2016). This therefore leads to variability of common sources of water in the region for successful aquaculture. It therefore became imperative to ascertain the physicochemical properties of commonly available sources of water in the area to ascertain its suitability for fish breeding. As posited by Bhatnagar and Devi, (2019) the determination of physicochemical characteristics of water is essential for assessing the suitability of water for various purposes like drinking, domestic, aquaculture and irrigation.

For that reason, a complete study on the physicochemical properties of common sources of water available to local fish farmers in Abakaliki is imperative to

determine its suitability for breeding operations. It is thus believed that the result of this research work will help solve the perennial problem of breeding failures and boost the production of *C. gariepinus* fingerlings particularly in Ebonyi state and Nigeria generally.

## **MATERIALS AND METHODS**

### **Study Area**

The research was conducted at the Hatchery unit, Department of Fisheries and Aquaculture Fish farm, Ebonyi State University, Abakaliki, Ebonyi State, Nigeria. Ebonyi State is situated between latitudes 6.24 °N and 6.28 °N and longitudes 7.00 °E and 7.06 °E on the South-East of Nigeria. The mean temperature range is 27°C to 31°C minimum and maximum respectively.

### **Brood Fish Selection and Acclimation**

Eight (4 males and 4 females) sexually matured samples of *C. gariepinus* brood stock size ranging from 800g-1.3kg total body weight was procured from a reputable fish farm in Abakaliki, Ebonyi State. All selected brood fish were transported to hatchery unit, Department of Fisheries and Aquaculture Fish farm, Ebonyi State University, Abakaliki in aerated tanks. Selection of gravid fish was done following the methods of (Gikonyo *et al.*, 2017). They

were kept separately in two 45 L indoor concrete tanks (2 m × 2 m × 1 m) and fed 35% crude protein diet (Madu, 2006) for two weeks before commencement of experiment.

### **Water Physicochemical Parameters**

The physicochemical parameters of the incubation water treatments were analyzed before the commencement and within the period of experiment. Mercury in glass thermometer was used to measure daily room and water temperature. The water pH was measured using the pH Hannah Portable Meter model HI 991 300. Dissolved oxygen (DO) of water was also measured using Hannah Portable Meter model HI 9142 and also by the modified Winkler azide method (APHA, 1989). Hannah Portable Meter model HI 9142 was equally used to determine water conductivity and total hardness. Nitrate, nitrite and ammonia in water were measured daily with the aid of Fresh Innovative Multitech, (NIFFRI) water testing kits.

### **Experimental Design**

The complete randomized design (CRD) was used for this experiment. The study adopted the partial water flow through incubation technique and lasted for approximately two months. Four different water sources

constituted the incubation media. They are: rain water (T1), borehole water (T2), river water (T3) and well water (T4). The incubation trough consisted of 4 hatchery circular 30 L plastic tanks. The circular 30 L plastic tanks were filled with the four different water media enumerated above and hereafter called treatments. Three replicates per water source at each trial were carried out to reduce sampling error. This brings a total of 12 circular 30 L plastic tanks. Prior to incubation, the tanks were properly washed with water and salt then dried. Each tank was securely laced with suspended 0.2mm nylon mesh net (kakabans) which will act as adhesion substrate for the eggs.

#### **Extraction of Milt, Stripping, Artificial Fertilization and Incubation of Eggs**

Extraction of milt was done following standard methods as described by Watson and Chapman (2002). The milt obtained from the gonad was squeezed into a beaker containing the normal saline water (0.9% chloride). After 9 hours' latency period at a room temperature of 27 °C and water temperature of 28 °C, the eggs were stripped. Egg stripping was done according to the method described by Amoah *et al.* (2020). Fertilization of eggs in this experiment was conducted following the procedure adopted

by Ndimele and Owodeinde (2012). This process facilitated a complete fertilization process and the fertilized eggs ready for incubation. With the aid of a sensitive weighing balance, (JT210N Electronic Top Loading weighing balance), 1 g of the fertilized eggs was measured, counted and gently spread, separated and well distributed on the pre- treated netting material inside the incubation tanks for each triplicate treatment. This procedure was done for all the three treatments and the entire set up was kept undisturbed in the hatchery for the period of the hatching session.

#### **Hatching and Percentage Hatchability of Fertilized Eggs' Estimation**

Incubated eggs were monitored and water temperature was maintained at 28° C. Larval movement was noticed between 18-24 hours. At the end of the hatching, duration of egg incubation for each water treatment was established. Dead, un-hatched eggs, hatched but dead larvae and hatched but deformed larvae were visually determined, counted and recorded. The netting was carefully removed and egg shells, dead larvae and all hatching remnants were siphoned out of the incubation media (Amoah *et al.*, 2021). The number of successfully hatched larvae were noted and recorded. Daily larval mortalities were

recorded for each water treatment. The percentage (%) hatchability was determined subjectively after 12-15 hours of fertilization by identifying the healthy developing eggs which were transparent and brownish in colour, while the dead eggs were estimated according to Abolude *et al.* (2013). Total number of successfully hatched larvae in each of the incubation water treatment and the number of deformed larvae were determined visually by simple count. The percentage hatchability was calculated according to the method described by Ndimele and Owodeinde (2012) using the formula:

$$\text{Hatchability (\%)} = \frac{\text{Total number of hatched eggs}}{\text{Total no of fertilized eggs}} \times 100$$

### **Larvae Management and Survival Percentage Estimation after 14-Days Culturing**

Feeding of the larvae commenced after three days when the yolk was absorbed. The hatchlings were fed *ad libitum* four times daily as follows: 9:00 am, 1:00 pm, 5:00 pm, and 9:00 pm with commercially prepared de-capsulated *Artemia nauplii* for 11 days. Leftover food particles, fry metabolites and other remnants were siphoned out of the

rearing tanks on daily basis before feeding. About 50% of water in each incubation tank was replaced with fresh water treatments accordingly so as to prevent the buildup of ammonia and the bowls carefully mopped with soft foam in order to remove dirt from the medium. Total renewal and washing of bowls with salt was done weekly (Abdulraheem *et al.*, 2020). At the end of 14 days' post hatching, the number of survived hatchlings in each water treatment was calculated and recorded. The percentage fry survivals up till the 14th day were estimated according to the method of Amachree *et al.* (2018) thus:

$$\text{Survival rate (\%)} = \frac{\text{Number of hatchlings alive up to larval stage on 14th day}}{\text{Total number of viable larvae initially hatched}} \times 100$$

### **Statistical Analysis**

Water physicochemical parameters such as temperature, dissolved oxygen, pH, ammonia, conductivity and total hardness were assessed for each individual treatment. Hatchability rate and survival rate were calculated for each individual treatment. Data from each treatment were then subjected to one-way analysis of variance (ANOVA) at  $p > 0.05$  significance level. The significance of


difference between means was determined by Duncan's Multiple Range Tests (DMRT) using the SPSS computer statistic package for windows 7 (version 21). Graphs were generated by using Microsoft Excel (2021) and all values were expressed as means  $\pm$  SE.

## RESULTS

Results of the mean water quality parameters tested during the 14 days breeding performance of *C. gariepinus* larvae in different water sources are presented in Table 2. There was no significant difference ( $P > 0.05$ ) in temperature, dissolved oxygen (DO) and ammonia values obtained during the experiment. The mean water temperature during the time of the experiment was ( $27.00 \pm 0.00^\circ\text{C}$  -  $28.00 \pm 0.00^\circ\text{C}$ ), while the mean


pH of water ranged between  $6.5 \pm 0.0$  -  $7.5 \pm 0.0$ . The mean dissolved oxygen value of water ranged between  $5.28 \pm 0.00$  -  $8.20 \pm 0.00$   $\text{mg}^{\text{l}}$  while mean values of ammonia ranged between  $0.21 \pm 0.00$  -  $0.40 \pm 0.00$   $\text{mg}^{\text{l}}$ .

However, there was a significant difference ( $P < 0.05$ ) in values obtained for conductivity and total hardness tests (Figure 1 and 2). Rain water (T1) recorded the lowest conductivity ( $24.3 \pm 0.88$   $\mu\text{S}^{\text{cm}}$ ) while T2 recorded the highest conductivity ( $383 \pm 0.00$   $\mu\text{S}^{\text{cm}}$ ) followed by T4 ( $285.2 \pm 0.00$   $\mu\text{S}^{\text{cm}}$ ), while T3 recorded ( $287.1 \pm 0.00$   $\mu\text{S}^{\text{cm}}$ ). Total hardness value was lowest in T1 ( $3.33 \pm 0.33$   $\text{mg}^{\text{l}}$ ) and highest in T2 ( $57.7 \pm 0.88$   $\text{mg}^{\text{l}}$ ) while T3 and T4 recorded  $32.3 \pm 0.33$   $\text{mg}^{\text{l}}$  and  $38.6 \pm 0.67$   $\text{mg}^{\text{l}}$  respectively.



**Figure 1: Mean Conductivity Values of Water Sources Used to Breed and Rear *C. gariepinus* fry.**


Vertical bars symbolize the mean  $\pm$  S.E. (Duncan's Multiple Range Test,  $P < 0.05$ ). Different letters indicate significant differences. \*T1=rain water, T2=borehole water, T3=river water, T4=well water



**Figure 2: Mean Total Hardness Values of Water Sources Used to Breed *C. gariepinus* fry.** Vertical bars symbolize the mean  $\pm$  S.E. (Duncan's Multiple Range Test,  $P < 0.05$ ). Different letters indicates significant differences. \*T1=rain water, T2=borehole water, T3=river water, T4=well water

### Breeding Performance


The results of the mean hatchability rates of *C. gariepinus* incubated in rearing units of different water media is presented in Table 3. There was significant difference ( $P > 0.05$ ) in hatching rates among the treatments. The mean values of percentage hatchability from the fertilized eggs incubated shows that hatchlings had relatively high mean values of (93.1 %) in T1, (92.0 %) in T4, 91.0 % in T3 and T2 obtained the least value of (84.9 %).



**Figure 3: Mean Hatchability Rates (%) of *C. gariepinus* eggs incubated in different Water Sources.** Vertical bars symbolize the mean  $\pm$  S.E. (Duncan's Multiple Range Test,  $P > 0.05$ ). Different letters indicate significant differences. \*T1=rain water, T2=borehole water, T3=river water, T4=well water.

### Survival Rates

The result of mean survival rates of *C. gariepinus* fry bred and reared for 14 days in different water media are shown in Figure 4. In the present study, larval survival rate was significantly different ( $P < 0.05$ ) in all four treatments. However, T2 recorded the highest larval survival rate representing 63.7 %, followed by T3 which recorded 61.4 % while T4 recorded 55.8 % and T1 recorded the lowest larval survival rate of 44.2 %.



**Figure 4: Mean Survival Rates (%) of *C. Gariepinus* Larvae Bred in different Water Sources. Vertical bars symbolize the mean  $\pm$  S.E. (Duncan's Multiple Range Test,  $P < 0.05$ ). Different letters indicate significant differences. \*T1=rain water, T2=borehole water, T3=river water, T4=well water**

## DISCUSSION

Water physicochemical parameters are critical in the biology and physiology of aquatic organisms (Amachree *et al.*, 2018). Table 1 shows acceptable range limits of vital water physicochemical parameters for the propagation of African catfishes. In this study, mean values obtained for temperature ( $^{\circ}\text{C}$ ) ( $27.00 \pm 0.00$   $^{\circ}\text{C}$  -  $28.00 \pm 0.00$   $^{\circ}\text{C}$ ), dissolved oxygen ( $5.28 \pm 0.00$  -  $8.20 \pm 0.00$   $\text{mg/l}$ ), pH ( $6.5 \pm 0.0$  -  $7.5 \pm 0.0$ ), and ammonia ( $0.21 \pm 0.00$  -  $0.40 \pm 0.00$   $\text{mg/l}$ ) were within

acceptable limits for successful propagation and culture of *C. gariepinus* (Table 2). These results indicate that these parameters did not negatively affect hatchability and survival of *C. gariepinus* larvae in all the treatments. This result is in agreement with the values obtained by Ukwe and Abu (2016); Alabi and Ocholi, (2019); Uka and Kalu (2019) in similar experiments.

However, rain water (T1) recorded a low conductivity value of  $24.3 \pm 0.88$   $\mu\text{S/cm}$  but values obtained for borehole water (T2), river water (T3) and well water (T4) were within acceptable range for the successful propagation of African catfish (Figure 1). According to Russell *et al.* (2011) water conductivity range of 200 - 500  $\mu\text{S/cm}$  is ideal for fish breeding. The low conductivity value recorded for T1 is not within the acceptable optimal conductivity range for fish propagation and rearing. This may have been responsible for the high larval mortalities and low survival rates observed in fries bred in T1. This finding aligns with the report of Makori *et al.* (2017) who assert that mid range conductivity of 200 - 1000  $\mu\text{S/cm}$  is favourable for larval development and fry survival but low conductivity of 0 - 200  $\mu\text{S/cm}$  and high conductivity of 1000 -



10,000  $\mu\text{S}/\text{cm}$  discourages larvae/fry survival.

In the same manner, rain water (T1) recorded a low total hardness value ( $3.33 \pm 0.33 \text{ mg}^{\text{l}}$ ) while the other three treatments recorded values that are within acceptable range for successful fish breeding (Figure 2). Krishnakumar *et al.* (2020) opined that the most excellent range of water hardness for optimal egg hatching and larvae viability of *C. gariepinus* is 30 – 500  $\text{mg}^{\text{l}} \text{CaCO}_3$ . So, the mortalities and poor growth performance recorded for larvae in T1 may be attributed to low water hardness. This finding aligns with the report of Silva *et al.* (2003) who asserted that low water hardness lowers hatching rates and reduces post hatch survival.

Hatchability rates were relatively high in all treatments but borehole water (T2) was significantly higher than others (Figure 3). These favourable hatching rates indicate that physicochemical parameters of the different water media may not have affected hatchability. Rather, it may be attributed to viability of the brood stock eggs and milt. Since the brood stocks are farm bred, they were fed ad libitum with feed containing nutrients in their correct proportions, thus improving their development and maturity (Amoah, *et al.*, 2020). Consequently, this

enabled the brood stocks to produce good quality eggs and milt in adequate quantity. The percentage hatchability rate recorded in the present study is similar to the results obtained by Olaniyi *et al.* (2018); Uka and Kalu (2019) and Amoah, *et al.* (2020).

Borehole water (T2) had the highest survival rate followed by well water (T4) and river water (T3) while rain water (T1) obtained the lowest value (Figure 4). The low larval survival rate observed in T1 may be as a result of variations in water physicochemical characteristics. Two physicochemical parameters namely: conductivity and total hardness were significantly low in T1 but were within acceptable range in other treatments. Conductivity and total hardness are two important parameters that affect larval and fry survival (Bhatnagar and Devi, 2019). Therefore, low conductivity and very low total hardness may have contributed to low survival rates of larvae bred and nursed in rain water treatment (T1). The survival rate recorded in the present study is similar to the results obtained by Olaniyi *et al.* (2018); Uka and Kalu (2019) in related experiments.

## CONCLUSION

Result from the study indicates that rain water may have the most suitable physicochemical qualities for incubating and

breeding *C. gariepinus* eggs but may not be suitable for larval, early fry culture. The study has equally demonstrated that river water, borehole water and well water have desirable physicochemical characteristics for breeding and culturing *C. gariepinus* larvae though borehole water may be the most suitable.

## REFERENCES

- Abolude, D. S., Opabunmi, O. O., Davies, O. A. and Awotoye, O. E. (2013). Study of Induced Breeding in *Clarias gariepinus* (Burchell, 1822) Using Ovaprim Hormone at Miracle Fish Farm, Zaria, Kaduna State, Nigeria. *Biological and Environmental Sciences Journal for the Tropics*, 10(4) .25-28.
- Akankali, J. A., Seiyaboh, E. I. and Abowei, J. F. N. (2011). Fish Hatchery Management in Nigeria. *Advanced Journal of Food Science and Technology* 3(2): 144-154.
- Akinwale A. O. and Faturoti, E. O. (2006). Biological performance of African Catfish (*Clarias gariepinus*) cultured in recirculating system in Ibadan, *Aquacultural Engineering* 36(1):18-23.DOI:10.1016/j.aquaeng.2006.05.001
- Akombo, P. M., Atile, J. I. and Obaje, J. A. (2018). The Effects of Some Physico-Chemical Parameters on the Breeding of Catfish (*Clarias gariepinus*) using Ovaprim and Ovatide. *International Journal of Innovative Studies in Aquatic Biology and Fisheries* 4(3): 1-9.  
DOI:http://dx.doi.org/10.20431/2454-7670.0403001 www.arcjournals.org
- Alabi, K. I. and Ocholi, O. S. (2019). Studies on the hatchability, growth performance and survival of African catfish (*Clarias gariepinus*) in different aquaculture production systems, *International Journal of Fisheries and Aquatic Studies* 7(5): 34-38.
- Amachree, D., Akinrotimi, O. A. and Dabor, P. O. (2018). Effects of Different Physiological Saline Concentrations on the Reproductive Performance of *Clarias gariepinus* (Burchell, 1822), *Journal of Fisheries Sciences* 12(4), 004-008.
- Amoah, K., Adu-Asiamah, P., Dong, X., Ampofo-Yeboah, K. and Abarike, E. D. (2020). A Comparative Study on the Hatchability and Survival Rate of African Catfish, *Clarias gariepinus* (Burchell, 1822), Induced with

- Catfish's Pituitary Gland Hormone from Farmed and Wild Sources. *Aquaculture International* <https://doi.org/10.1007/S10499-020-00591-4>.
- Ataguba, G. A., Okomoda, V. T. and Azave, V. T. (2015). Effect of Photoperiod on Hatching Success of Eggs and Early Fry Performance of *Clarias gariepinus* (Siluriformes, Burchell 1822), *Trakia Journal of Sciences* 2: 171-174, Doi:10.15547/Tjs.2015.02.010
- Bhantanagar, A. and Devi, P. (2019). Water Quality Guidelines for Fish Management of Pond Fish Culture. *International Journal of Environmental Sciences* 3(6): 1-30.
- Ebiriekwe, S. C. (2012). Studies on the Suitability of Different Sources of Water of Abakiliki Municipality for the Propagation of *Clarias gariepinus* (Burchell 1822) “(Unpublished Master's Dissertation)”. Ebonyi State University, Abakaliki.
- Gikonyo, J. M., Kibegwa, F. M. and Inyangala, B. O. (2017). Comparative Assessment of Hatching Rates of African Catfish (*Clarias gariepinus*) Eggs Using Nile Cabbage and Kaka Bans Substrates. *Livestock Research for Rural Development* 29(164).
- Kareem, O. K., Ajani, E. K., Akintunde, M. A., Olanrewaju, A. N. and Oduntan, O. B. (2017). Effect of Different Fertilization and Egg De-adhesion Methods on Hatching and Survival of *Clarias gariepinus* (Burchell 1822) Fry, *Journal of Fisheries Sciences* 11(1): 021-027
- Krishnakumar, A., Anton, E. S. P. and Jayawardena, U. A. (2020). Water hardness influenced variations in reproductive potential of two freshwater fish species; *Poecilia reticulata* and *Betta splendens*. *BMC Research Notes* 13, 542-548 <https://doi.org/10.1186/s13104-020-05382-x>
- Madu, C. T. (2006). The Effects of Brood Stock Size on the Economy of Catfish (*Clarias anguillaris*) Fry Production Using the Hormone Induced Natural Breeding Technique. *Journal of Aquatic Science* 21: 19-22.
- Makori, A. J., Abuom, P. O., Kapiyo, R., Anyona, D. N. and Dida, G. O. (2017). Effects of water physicochemical parameters on tilapia (*Oreochromis niloticus*) growth in earthen ponds in Teso North Sub-

- County, Busia County. *Fisheries and Aquatic Sciences* 20, 30-40. DOI 10.1186/s41240-017-0075-
- Ndimele, P. E. and Owodeinde, F. G. (2012). Comparative Reproductive and Growth Performance of *Clarias gariepinus* (Burchell, 1822) and Its Hybrid Induced with Synthetic Hormone and Pituitary Gland of *Clarias gariepinus*. *Turkish Journal of Fisheries and Aquatic Sciences*, 12. Pp.619-626. DOI: 10.4194/1303-2712-v12\_3\_09
- Nwabunike, M. O. (2016). Possible Changes in Heavy metals Bioaccumulation in Fish Liver in selected Rivers of Ebonyi state. *Nigerian Journal of Biotechnology*, 31: 66 –73. DOI: <http://dx.doi.org/10.4314/njb.v31i1.9>
- Nwakanma, C. and Okwum, C. (2015). The Evaluation of Groundwater Quality on Successful Hatchery Culture of African Catfish (*Clarias gariepinus*) In Ikwuano Oboro, Abia State. *World Journal of Pharmaceutical Research* 5(1): 12-23.
- Olaniyi, C. O., Alalade, E. O., Shittu, R. O., Adesina, O. R., Osungbade, K. D., Adelokun, A. M. (2018). Effects of Different Water Sources on Hatchability, Survival and Growth Performance of African Catfish (*Clarias gariepinus*), *International Journal of Innovative Research and Advanced Studies* 5: 5.
- Onada, O. A. and Ogunola, O. S. (2017). Effects of Catfish (*Clarias gariepinus*) Brood-stocks Egg Combination on Hatchability and Survival of Fish Larvae, *Journal of Aquaculture Research and Development* Doi:10.4172/2155-9546.S2-014
- Opoke, S. U. and Osayande, A. D. (2018). Physicochemical, Major Anions and Heavy Metals in Surface Water and Groundwater at Pb–Zn Mining areas of Ebonyi State, *Nigeria Journal of Applied Science and Environmental Management*, 22(7): 1017-1020. DOI: <https://dx.doi.org/10.4314/jasem.v22i7.3>
- Russell, M., Shuke, R. and Samantha, S. (2011). Effects of Conductivity on Survivorship and Weight of Goldfish (*Carassius auratus*). Available at <http://departments.juniata.edu/biology/eco/documents/Russell>.
- Silva, L. V. F., Golombieski, J. I. and Baldisserotto, B. (2003). Incubation of Silver Catfish, *Rhamdia quelen*

- (Pimelodidae), Eggs at Different Calcium and Magnesium Concentrations. *Aquaculture*, 228: 279–287.
- Summerfelt, R. C. (1998). Water quality considerations for aquaculture. Wiley & Sons, Inc., New York. Pp. 23.
- Uka, A. and Kalu, S. E. (2019). Differences in physiochemical properties of water from neighborhood boreholes and their usefulness in *Clarias gariepinus* egg hatching. *Journal of Applied Science and Environmental Management* 28(8): 1567-1571 Doi: <https://dx.doi.org/10.4313/jasem.v2318.22>.
- Ukwe, I. O. K. and Abu, O. M. G. (2016). Physico-Chemical Parameters of Water in Holding Tanks of *Clarias gariepinus* Induced with Ovaprim and Ovulin Hormones. *International Journal of Innovative Studies in Aquatic Biology and Fisheries* 2(4); 12-19. <http://dx.doi.org/10.20431/2454-7670.0204003> [www.arcjournals.org](http://www.arcjournals.org)

**Table 1: Water Quality Requirements of African Catfish Hatchery**

Parameters	Acceptable Range Limits
Temperature	26-28 °C
Dissolved Oxygen (DO)	>4 mgL <sup>-1</sup>
Water pH	7-7.5
Ammonia	0-0.5 mg/l
Conductivity	200–1000 µS/cm
Total Hardness	30 – 180 mg/l

Source: Pronob Das *et al.*, (2011).

**Table 2: Mean Physicochemical Parameters of Different Water Sources Used to Breed and Culture *Clarias gariepinus* Larvae for 14 days.\***

Parameters	Treatments			
	T1	T2	T3	T4
Temperature (°C)	27.00±0.00 <sup>a</sup>	28.00±0.00 <sup>a</sup>	27.00±0.00 <sup>a</sup>	28.00±0.00 <sup>a</sup>
Dissolved Oxygen (mg/l)	6.20±0.00 <sup>a</sup>	5.82±0.00 <sup>a</sup>	7.20±0.00 <sup>a</sup>	8.20±0.00 <sup>a</sup>
pH	7.00±0.00 <sup>a</sup>	6.50±0.00 <sup>a</sup>	7.50±0.00 <sup>a</sup>	6.50±0.00 <sup>a</sup>
Ammonia (mg/l)	0.21±0.00 <sup>a</sup>	0.40±0.00 <sup>a</sup>	0.21±0.00 <sup>a</sup>	0.38±0.00 <sup>a</sup>
Conductivity (µS/cm)	24.3±0.88 <sup>a</sup>	383.3±0.78 <sup>b</sup>	285.2±0.00 <sup>c</sup>	287.1±0.00 <sup>b</sup>
Total Hardness (mg/l)	3.33±0.33 <sup>a</sup>	57.7±0.88 <sup>c</sup>	32.3±0.33 <sup>b</sup>	38.6±0.67 <sup>b</sup>

\*Values in the same row with different superscripts are significantly different (P < 0.05)

\*T1= rain water, T2=borehole water, T3=river water, T4=well water

**Table 3: Hatchability/Survival Rates of *Clarias gariepinus* Cultured in Different Water Media for 14 days\***

Parameters				Treatments			
				T1	T2	T3	T4
Estimated	Number	of	Eggs	$640 \pm 0.00^a$	$640 \pm 0.00^a$	$640 \pm 0.00^a$	$640 \pm 0.00^a$
used/treatment							
Estimated Number of Fertilized Eggs				$354 \pm 12.5^a$	$344 \pm 10.9^a$	$367 \pm 10.9^a$	$355 \pm 5.56^a$
Estimated Number of Eggs Hatched				$333.0 \pm 15.2^a$	$292.0 \pm 3.21^b$	$322.0 \pm 3.21^b$	$326.7 \pm 6.6^a$
Hatchability Rate (%)				$93.1 \pm 1.60^b$	$84.9 \pm 3.12^a$	$91.0 \pm 3.12^{ab}$	$92.0 \pm 2.21^b$
Larval Survival Rate at 14 days (%)				$44.2 \pm 0.55^a$	$63.7 \pm 2.41^c$	$55.8 \pm 3.00^b$	$61.4 \pm 3.00^{bc}$

\*Values in the same row with different superscripts are significantly different ( $P < 0.05$ )

\*T1= Rain water, T2=Borehole water, T3=River water, T4=Well water